

Embedding of C_q and R_q into noncommutative L_p -spaces, $1 \leq p < q \leq 2$

Quanhua Xu

Abstract

We prove that a quotient of subspace of $C_p \oplus_p R_p$ ($1 \leq p < 2$) embeds completely isomorphically into a noncommutative L_p -space, where C_p and R_p are respectively the p -column and p -row Hilbertian operator spaces. We also represent C_q and R_q ($p < q \leq 2$) as quotients of subspaces of $C_p \oplus_p R_p$. Consequently, C_q and R_q embed completely isomorphically into a noncommutative $L_p(M)$. We further show that the underlying von Neumann algebra M cannot be semifinite.

0 Introduction

In the recent remarkable work [J1], Junge proved that Pisier's space OH (the operator Hilbert space) embeds completely isomorphically into the predual of a QWEP von Neumann algebra. He further showed in [J2] that the von Neumann algebra in question can be chosen to be hyperfinite. Junge's proof consists of two steps. The first one is to represent OH (in a precise way) as a quotient of subspace of $C \oplus R$, where C and R are the usual column and row Hilbertian operator spaces. (By a quotient of subspace of X we mean a quotient of a subspace Y of X .) His tool for this representation theorem is a formula due to Pusz and Woronowicz on the square root of the product of two commuting positive operators on a Hilbert space (cf. [PW]). The second step is a kind of Khintchine type inequality involving a three term K-functional. Junge obtained such an inequality via reduced free product by using Speicher's central limit theorem (cf. [Sp]).

Junge's arguments were simplified by Pisier [P4]. The fact that OH is a quotient of subspace of $C \oplus R$ has been already known before [J1] (see [P1, Exercise 7.9]). Pisier's approach to this is via the complex interpolation. Junge's second step above was then replaced in [P4] by a Khintchine type inequality for the generalized circular systems obtained in [PS].

The fact that the complex interpolation can be used instead of Pusz-Woronowicz's formula is not surprising since this formula is intimately related to interpolation. Indeed, it can be shown that Pusz-Woronowicz's formula is equivalent to the real interpolation of weighted L_2 -spaces. This will be discussed in details in the forthcoming work [JX3] (see also section 3 below for a brief discussion).

The purpose of the present paper is to consider the same embedding problem for the p -column and p -row spaces C_p and R_p . Recall that C_p (resp. R_p) is the subspace of the Schatten class S_p consisting of matrices whose all entries but those in the first column (resp. row) vanish. Both C_p and R_p are isometric to ℓ_2 as Banach spaces. Note that C_2 and R_2 coincide with OH (completely isometrically), and C_∞ and R_∞ are just C and R respectively.

We will prove that if $1 \leq p < q \leq 2$, then both C_q and R_q embed completely isomorphically into a noncommutative L_p -space. An intermediate result is that C_q and R_q are quotients of subspace

Laboratoire de Mathématiques, Université de Franche-Comté, 25030 Besançon, cedex - France
 qx@math.univ-fcomte.fr

2000 *Mathematics subject classification*: Primary 46L07; Secondary 47L25

Key words and phrases: embedding, p -column and p -row spaces, noncommutative L_p -spaces, interpolation

of $C_p \oplus_p R_p$. We will present in section 3 two proofs for this representation result. The first one is via the complex interpolation as in [P4], whereas the second is based on the real interpolation.

Besides the representation of C_q as a quotient of subspace of $C_p \oplus_p R_p$, another ingredient of the embedding theorem above is the Khintchine type inequality for generalized circular systems in noncommutative L_p -spaces from [X1]. We will recall in section 4 this inequality, and also give its Fermionic analogue (at least for $p > 1$).

More generally, we will show that any quotient of subspace of $C_p \oplus_p R_p$ ($1 \leq p < 2$) embeds completely isomorphically into a noncommutative L_p . This will be done by the Khintchine inequality quoted previously and a decomposition result for subspaces of $C_p \oplus_p R_p$ (see Proposition 2.1). It was proved in [X1] that if an operator space X and its dual embed completely isomorphically into a noncommutative L_p ($1 < p < 2$), then X is completely isomorphic to a quotient of subspace of $H_p^c \oplus_p K_p^r$ for two Hilbert spaces H and K , where H_p^c (resp. K_p^r) means that H (resp. K) is equipped with the p -column (resp. p -row) operator space structure. Therefore, we obtain that for $1 < p < 2$ an operator space X is completely isomorphic to a quotient of subspace of $H_p^c \oplus_p K_p^r$ iff both X and X^* embed completely isomorphically into a noncommutative L_p .

To end this introduction let us mention that all algebras constructed in [J1, J2, P4] are of type III. This is not a hazard for Pisier [P5] proved that OH cannot embed completely isomorphically into the predual of a semifinite von Neumann algebra. We will also extend this result by Pisier to our general setting: C_q cannot embed completely isomorphically into a semifinite L_p ($1 \leq p < q \leq 2$).

1 Preliminaries

In this section we recall briefly the natural operator space structure on noncommutative L_p -spaces and some elementary properties of column and row spaces. We use standard notions and notation from operator space theory (cf. [ER, P1]).

It is well known that there are several equivalent constructions of noncommutative L_p -spaces associated with a von Neumann algebra. All these constructions give the same spaces (up to isometry). In this paper we will use Haagerup noncommutative L_p -spaces constructed in [H]. [Te1] gives an excellent exposition of the subject. Let M be a von Neumann algebra and $1 \leq p \leq \infty$. As usual, $L_p(M)$ denotes the Haagerup L_p -space associated with M . Recall that $L_\infty(M)$ is just M itself, and $L_1(M)$ can be identified with the predual M_* of M . More generally, if $1 \leq p < \infty$ and $1/p + 1/p' = 1$, the dual of $L_p(M)$ is $L_{p'}(M)$. When M is semifinite, we will always consider $L_p(M)$ as the usual L_p -space constructed from a normal semifinite faithful trace.

Now we describe the natural operator space structure on $L_p(M)$ as introduced in [P2, P1]. For $p = \infty$, $L_p(M) = M$ has its natural operator space structure as a von Neumann algebra. For $p = 1$, the natural operator space structure on $L_1(M) = M_*$ is induced by the natural embedding of M_* into its bidual M^* , the latter being the standard dual of M . In fact, as explained in [P1, section 7], it is more convenient to consider $L_1(M)$ as the predual of the opposite von Neumann algebra M^{op} , which is isometric (but in general not completely isomorphic) to M , and to equip $L_1(M)$ with the operator space structure inherited from $(M^{\text{op}})^*$. The main reason for this choice is that the duality between S_∞^n and S_1^n in operator space theory is the parallel duality (instead of the usual tracial duality). Note that this choice further insures that the equality $L_1(M_n \otimes M) = S_1^n \hat{\otimes} L_1(M)$ (operator space projective tensor product) holds true. Finally, the operator space structure on $L_p(M)$ is obtained by complex interpolation, using the well known interpretation of $L_p(M)$ as the interpolation space $(M, L_1(M))_{1/p}$ (see [Te2]). If M admits a normal faithful state φ , we can also use Kosaki's interpolation [Ko].

In particular, the Schatten classes S_p are equipped with their natural operator space structure. We will need vector-valued Schatten classes as introduced in [P2]. Let E be an operator space. $S_1[E]$ is defined as the operator space projective tensor product $S_1 \hat{\otimes} E$. Then for any $1 < p < \infty$, $S_p[E]$ is defined by interpolation:

$$S_p[E] = (S_\infty[E], S_1[E])_{1/p}.$$

When E is a subspace of a noncommutative $L_p(M)$, the norm of $S_p[E]$ admits a very simple description. Indeed, $S_p[L_p(M)]$ coincides (completely isometrically) with $L_p(B(\ell_2) \overline{\otimes} M)$ ($p < \infty$). If $E \subset L_p(M)$, then $S_p[E]$ is the closure in $L_p(B(\ell_2) \overline{\otimes} M)$ of the algebraic tensor product $S_p \otimes E$.

By reiteration, for any $1 \leq p_0, p_1 \leq \infty$ and $0 < \theta < 1$

$$(S_{p_0}[E], S_{p_1}[E])_\theta = S_p[E],$$

where $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$. More generally, given a compatible couple (E_0, E_1) of operator spaces we have

$$(1.1) \quad (S_{p_0}[E_0], S_{p_1}[E_1])_\theta = S_p[(E_0, E_1)_\theta].$$

The main concern of this paper is on the L_p -space counterparts of the usual row and column spaces R and C . Let C_p (resp. R_p) denote the subspace of S_p consisting of matrices whose all entries but those in the first column (resp. row) vanish. It is clear that C_p and R_p are completely 1-complemented subspaces of S_p . We have the following completely isometric identifications:

$$(1.2) \quad (C_p)^* \cong C_{p'} \cong R_p \quad \text{and} \quad (R_p)^* \cong R_{p'} \cong C_p, \quad \forall 1 \leq p \leq \infty.$$

C_p and R_p can be also defined via interpolation from C and R . We view (C, R) as a compatible couple by identifying both of them with ℓ_2 (at the Banach space level), i.e. by identifying the canonical bases $(e_{k,1})$ of C_p and $(e_{1,k})$ of R_p with (e_k) of ℓ_2 . Then

$$C_p = (C, R)_{1/p} = (C_\infty, C_1)_{1/p} \quad \text{and} \quad R_p = (R, C)_{1/p} = (R_\infty, R_1)_{1/p}.$$

By reiteration, for any $1 \leq p_0, p_1 \leq \infty$ and $0 < \theta < 1$

$$(1.3) \quad C_p = (C_{p_0}, C_{p_1})_\theta \quad \text{and} \quad R_p = (R_{p_0}, R_{p_1})_\theta,$$

where $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$. Like C and R , C_p and R_p are 1-homogenous 1-Hilbertian operator spaces. We refer to [P2, P3] for the proofs of all these elementary facts.

More generally, given a Hilbert space H and $1 \leq p \leq \infty$ we denote by H_p^c (resp. H_p^r) the Schatten p -class $S_p(\mathbb{C}, H)$ (resp. $S_p(\bar{H}, \mathbb{C})$) equipped with its natural operator space structure. H_p^c (resp. H_p^r) can be naturally viewed as the column (resp. row) subspace of the Schatten class $S_p(H)$ (resp. $S_p(\bar{H})$). When H is separable and infinite dimensional, H_p^c and H_p^r are respectively C_p and R_p above. All previous properties for C_p and R_p hold for H_p^c and H_p^r too. H_p^c and H_p^r are respectively the p -column and p -row spaces associated with H .

Now let E be an operator space. We denote by $C_p[E]$ (resp. $R_p[E]$) the closure of $C_p \otimes E$ (resp. $R_p \otimes E$) in $S_p[E]$. Again, $C_p[E]$ and $R_p[E]$ are completely 1-complemented subspaces of $S_p[E]$. If E is a subspace of a noncommutative $L_p(M)$, the norm of $C_p[E]$ is easy to be determined as follows. For any finite sequence $(x_k) \subset E$

$$\left\| \sum_k x_k \otimes e_k \right\|_{C_p[E]} = \left\| \left(\sum_k x_k^* x_k \right)^{1/2} \right\|_{L_p(M)},$$

where (e_k) denotes the canonical basis of C_p . More generally, if $a_k \in C_p$, then

$$(1.4) \quad \left\| \sum_k x_k \otimes a_k \right\|_{C_p[E]} = \left\| \left(\sum_{k,j} \langle a_k, a_j \rangle x_k^* x_j \right)^{1/2} \right\|_{L_p(M)},$$

where $\langle \cdot, \cdot \rangle$ denotes the scalar product in C_p . (In terms of matrix product, $\langle a_k, a_j \rangle = a_k^* a_j$.) We have also a similar description for the norm of $R_p[E]$. The same formulas hold for more general $H_p^c[E]$ and $H_p^r[E]$.

2 A decomposition

Let $T : X \rightarrow Y$ be a linear map between two Banach spaces. Recall that the graph (or domain) of T is the subspace of $X \oplus Y$ given by

$$G(T) = \{(x, Tx) : x \in \text{Dom}(T)\},$$

where $\text{Dom}(T)$ stands for the definition domain of T .

Proposition 2.1 *Let X_0, X_1 be 1-homogeneous 1-Hilbertian operator spaces. Let $X = X_0 \oplus_2 X_1$ and $Y \subset X$ be a closed subspace. Then there are closed subspaces $Y_0, Z_0 \subset X_0$, $Y_1, Z_1 \subset X_1$ and an injective closed densely defined operator T from Z_0 to Z_1 with dense range such that*

$$Y = Y_0 \oplus_2 Y_1 \oplus_2 G(T) \quad \text{with completely equivalent norms,}$$

where $G(T)$ is the graph of T regarded as a subspace of $Z_0 \oplus_2 Z_1 \subset X_0 \oplus_2 X_1$. Moreover, the relevant constants in the above equivalence are universal.

Proof. Let $\langle \cdot, \cdot \rangle_i$ denote the scalar product of X_i , and let P_i be the natural projection from X onto X_i ($i = 0, 1$). Since the map $(x, y) \mapsto \langle P_0(x), P_0(y) \rangle_0$ is a contractive sesquilinear form on Y , there is a contraction $A : Y \rightarrow Y$ such that

$$(2.1) \quad \langle x, A(y) \rangle = \langle P_0(x), P_0(y) \rangle_0, \quad x, y \in Y.$$

It is clear that $A \geq 0$ and

$$(2.2) \quad \langle x, (1 - A)(y) \rangle = \langle P_1(x), P_1(y) \rangle_1, \quad x, y \in Y.$$

Therefore,

$$x \in \ker A \Leftrightarrow P_0(x) = 0 \quad \text{and} \quad x \in \ker(1 - A) \Leftrightarrow P_1(x) = 0.$$

Set

$$Y_0 = \ker(1 - A) \quad \text{and} \quad Y_1 = \ker A.$$

Then it follows that

$$P_i|_{Y_i} = A|_{Y_i}, \quad i = 0, 1.$$

Consequently,

$$Y_i \subset X_i, \quad i = 0, 1.$$

Let Z be the orthogonal complement of $Y_0 \oplus_2 Y_1$ in Y . Then we have the following orthogonal decomposition of Y at the Banach space level:

$$(2.3) \quad Y = Y_0 \oplus_2 Y_1 \oplus_2 Z.$$

We will show that this decomposition also holds in the category of operator spaces with completely equivalent norms.

Let $y_0 \in Y_0$ and $z \in Z$. Then by (2.1) and the definition of Y_0

$$\langle y_0, P_0(z) \rangle_0 = \langle P_0(y_0), P_0(z) \rangle_0 = \langle y_0, A(z) \rangle = \langle A(y_0), z \rangle = \langle y_0, z \rangle = 0.$$

Therefore, Y_0 and $P_0(Z)$ are orthogonal subspaces of X_0 . Let Q_0 be the orthogonal projection from X_0 onto Y_0 . Since X_0 is homogeneous, Q_0 is completely contractive. Put $R_0 = Q_0 P_0|_Y$. Then R_0 is a completely contractive projection from Y onto Y_0 . In fact, it is easy to see that R_0 is exactly the orthogonal projection onto Y_0 . Indeed, we have $R_0 = 0$ on Y_1 , and as well as on Z for $Q_0|_{P_0(Z)} = 0$. Thus $R_0 = 0$ on the orthogonal complement of Y_0 . Similarly, the orthogonal

projection R_1 from Y on Y_1 is also completely contractive. It follows that the decomposition in (2.3) holds at the operator space level with completely equivalent norms.

It remains to express the third space Z as a graph. To that end let us observe that both A and $1 - A$ are injective operators on Z with dense range. Consequently, the inverses of A and $1 - A$ are densely defined operators on Z too.

On the other hand, since A and $P_0|_Y$ (resp. $1 - A$ and $P_1|_Y$) have the same kernel, we deduce that any $z \in Z$ is uniquely determined by $P_0(z)$ as well as by $P_1(z)$. Therefore, Z must be the graph of a closed densely defined operator, i.e. the graph of the map $P_0(z) \mapsto P_1(z)$. Let us explain this elementary fact precisely in the present situation. Let Z_i be the closure of $P_i(Z)$ in X_i . By (2.1) and (2.2), there are unitaries u_i from Z_i to Z such that

$$A^{1/2}|_Z = u_0 P_0|_Z \quad \text{and} \quad (1 - A)^{1/2}|_Z = u_1 P_1|_Z.$$

Therefore, for any $z \in Z$ we have

$$P_1(z) = u_1^*(1 - A)^{1/2}(z) = u_1^*(1 - A)^{1/2}A^{-1/2}u_0(P_0(z)).$$

Thus

$$T = u_1^*(1 - A)^{1/2}A^{-1/2}u_0$$

is a closed densely defined operator from Z_0 to Z_1 whose domain coincides with Z . Note that T is invertible and its inverse is also densely defined. This completes the proof. $\square$

Remark 2.2 Let $T = w\Delta$ be the polar decomposition of T . Then w is a unitary and Δ is a positive self-adjoint non-singular closed densely defined operator on Z_0 . By the homogeneity of X_1 and replacing Z_1 by Z_0 (but still equipped with the operator space structure of X_1), we can assume $T = \Delta$ is positive. Let us note, however, that with the previous precise form of T we can simply take $(1 - A)^{1/2}A^{-1/2}$ as Δ .

Applying the previous proposition to row and column spaces, we get the following

Corollary 2.3 *Let H, K be Hilbert spaces and $1 \leq p \leq \infty$. Let E be a closed subspace of $H_p^c \oplus_p K_p^r$. Then there are closed subspaces $H_0, H_1 \subset H$, $K_0, K_1 \subset K$ and an injective closed densely defined operator D from H_1 to K_1 with dense range such that*

$$E = (H_0)_p^c \oplus_p (K_0)_p^r \oplus_p G(D) \quad \text{with completely equivalent norms,}$$

where $G(D)$ is the graph of D regarded as a subspace of $(H_1)_p^c \oplus_p (K_1)_p^r$. Moreover, the relevant constants in the above equivalence are universal.

Remark. In the situation of the above corollary, it suffices to consider the case of $p \leq 2$ or $p \geq 2$ for we have

$$H_p^c \cong H_{p'}^r \quad \text{completely isometrically.}$$

3 C_q as a quotient of subspace of $C_p \oplus_p R_p$

It is known that OH can be represented as a quotient of subspace of $C \oplus R$ (see [P1, Exercise 7.9] and [P4]). Pisier's proof is via complex interpolation by using his interpolation formula $OH = (C, R)_{1/2}$ obtained in [P3]. The aim of this section is to show a similar result for C_q relative to the couple (C_p, R_p) for any q between p and p' (recalling that p' denotes the index conjugate to p).

Theorem 3.1 *Let $1 \leq p \leq \infty$ and $\min(p, p') < q < \max(p, p')$. Then C_q and R_q are completely isometric to quotients of subspace of $C_p \oplus_p R_p$.*

We will present two proofs for the theorem above. The first is based on the complex interpolation like in [P4]. The second one is similar to Junge's approach to the representation of OH as a quotient of subspace of $C \oplus_\infty R$ (cf. [J1]); but instead of Pusz-Woronowicz's formula, we will use directly the real interpolation. Both approaches have their advantages. The first one is more transparent and slightly simpler. Moreover, it shows that C_q is completely isometric to a quotient of subspace of $C_p \oplus_p R_p$. The major advantage of the second proof is that it gives the precise form of the quotient of subspace (or the two densities involved in this quotient) of $C_p \oplus_p R_p$ which is completely isomorphic to C_q . This precise form is crucial for many applications.

We will need the following lemma from [X1], which is a generalization of [P3, Theorem 8.4]. See also [X2] for a more general result.

Lemma 3.2 *Let $2 \leq p \leq \infty$, $0 < \theta < 1$, $\frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{p'}$ and r be the conjugate index of $p/2$. Let (e_k) denote the canonical basis of C_q . Then for any finite sequence $(x_k) \subset S_p$*

$$\left\| \sum_k x_k \otimes e_k \right\|_{S_p[C_q]} = \sup \left\{ \left(\sum_k \|\alpha x_k \beta\|_2^2 \right)^{1/2} \right\},$$

where the supremum runs over all α and β in the unit ball of $S_{2r\theta-1}$ and $S_{2r(1-\theta)-1}$, respectively. Moreover, the supremum can be restricted to all α and β in the positive parts of these unit balls.

Proof of Theorem 3.1. This proof is similar to the corresponding proof on the representation of OH as a quotient of subspace of $C \oplus R$ in [P4]. Since $R_q \cong C_{q'}$, it suffices to prove the assertion for C_q . By symmetry, we may assume $p \geq 2$. We will use the following interpolation formula (see (1.2) and (1.3))

$$(3.1) \quad C_q = (C_p, R_p)_\theta,$$

where θ is determined by $\frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{p'}$. To proceed further to the proof, we need to recall some elementary facts on complex interpolation. Let (X_0, X_1) be a compatible couple of Banach spaces. Set $S = \{z \in \mathbb{C} : 0 \leq \operatorname{Re}(z) \leq 1\}$. Let μ be the harmonic measure in S at the point θ . Note that μ can be written as $\mu = (1-\theta)\mu_0 + \theta\mu_1$, where μ_0 and μ_1 are probability measures respectively on the left boundary ∂_0 and the right boundary ∂_1 of S . Let $\mathcal{F}(X_0, X_1)$ be the family of all functions f from S to $X_0 + X_1$ satisfying the following conditions:

- i) f is analytic in the interior of S ;
- ii) $f|_{\partial_j} \in L_2(X_j; \mu_j)$, $j = 0, 1$.

$\mathcal{F}(X_0, X_1)$ is equipped with the norm

$$(3.2) \quad \|f\| = \left[(1-\theta) \|f|_{\partial_0}\|_{L_2(X_0; \mu_0)}^p + \theta \|f|_{\partial_1}\|_{L_2(X_1; \mu_1)}^p \right]^{\frac{1}{p}}$$

(with the obvious change for $p = \infty$). Then the complex interpolation space $(X_0, X_1)_\theta$ can be defined as the space of all $x \in X_0 + X_1$ such that there is $f \in \mathcal{F}(X_0, X_1)$ such that $f(\theta) = x$, equipped with the quotient norm

$$\|x\|_\theta = \inf \{ \|f\|_{\mathcal{F}(X_0, X_1)} : f(\theta) = x, f \in \mathcal{F}(X_0, X_1) \}.$$

We refer to [BL] for more details. We should call the reader's attention to the fact that the ℓ_p -norm in (3.2) is usually replaced by the ℓ_2 -norm (to be consistent with the L_2 -norm). However, at the end we get the same interpolation norm $\|\cdot\|_\theta$. (In fact, the indices 2, p in (3.2) can be replaced by any two indices in $[1, \infty]$ without changing the norm $\|\cdot\|_\theta$.) Note that $\mathcal{F}(X_0, X_1)$ is a subspace of

$$\mathcal{E}(X_0, X_1) = (1-\theta)^{\frac{1}{p}} L_2(X_0; \mu_0) \oplus_p \theta^{\frac{1}{p}} L_2(X_1; \mu_1).$$

Using the Poisson integration, we easily see that $\mathcal{E}(X_0, X_1)$ is exactly the space of all functions from S into $X_0 + X_1$ harmonic inside S and verifying the condition ii) above. Then $\mathcal{F}(X_0, X_1)$ is the subspace of $\mathcal{E}(X_0, X_1)$ consisting of analytic functions.

Now return back to the formula (3.1). In view of the above discussion set

$$H = (1 - \theta)^{\frac{1}{p}} L_2(\ell_2; \mu_0) \quad \text{and} \quad K = \theta^{\frac{1}{p}} L_2(\ell_2; \mu_1).$$

H and K are infinite dimensional separable Hilbert spaces. Let

$$E = H_p^c \oplus_p K_p^r.$$

As mentioned above, E can be considered as the space of ℓ_2 -valued harmonic functions in S with square integrable boundary values. Let $F \subset E$ be the subspace of analytic functions. Thus F is a subspace of $C_p \oplus_p R_p$. Consider the map $Q : F \rightarrow C_q$ defined by $Q(f) = f(\theta)$. We are going to show that Q is a completely isometric surjection. By virtue of [P2, Lemma 1.7], this is equivalent to show that $I_{S_p} \otimes Q$ extends to a quotient map from $S_p[F]$ to $S_p[C_q]$.

First we prove that $I_{S_p} \otimes Q$ extends to a contraction from $S_p[F]$ to $S_p[C_q]$ (which amounts to saying $\|Q\|_{cb} \leq 1$). Let $f \in S_p[F]$ be of norm ≤ 1 (assuming that f is a finite matrix). Let $x = f(\theta)$. We have to show $\|x\|_{S_p[C_q]} \leq 1$. To this end we use Lemma 3.2. Let f_n be the n^{th} coordinate of f . Set $x_n = f_n(\theta)$. Let α (resp. β) be a positive unit element in $S_{2r\theta^{-1}}$ (resp. $S_{2r(1-\theta)^{-1}}$). Define for $z \in S$

$$g(z) = \sum_n g_n(z) \otimes e_n \quad \text{with} \quad g_n(z) = \alpha^{\frac{z}{\theta}} f_n(z) \beta^{\frac{1-z}{1-\theta}}.$$

Then g_n is an analytic function in S with values in S_2 and

$$x_n = g_n(\theta) = (1 - \theta) \int_{\partial_0} g_n(z) d\mu_0(z) + \theta \int_{\partial_1} g_n(z) d\mu_1(z).$$

By the Hölder inequality, we have

$$\begin{aligned} \int_{\partial_0} \sum_n \|g_n(z)\|_2^2 d\mu_0(z) &\leq \int_{\partial_0} \sum_n \|f_n(z) \beta^{\frac{1}{1-\theta}}\|_2^2 d\mu_0(z) \\ &= \int_{\partial_0} \sum_n \text{Tr}(\beta^{\frac{1}{1-\theta}} f_n(z)^* f_n(z) \beta^{\frac{1}{1-\theta}}) d\mu_0(z) \\ &\leq \left\| \sum_n \int_{\partial_0} f_n(z)^* f_n(z) d\mu_0(z) \right\|_{p/2}. \end{aligned}$$

Similarly,

$$\int_{\partial_1} \sum_n \|g_n(z)\|_2^2 d\mu_1(z) \leq \left\| \sum_n \int_{\partial_1} f_n(z) f_n(z)^* d\mu_1(z) \right\|_{p/2}.$$

On the other hand, noting

$$S_p[E] = S_p[H_p^c] \oplus_p S_p[K_p^r] = H_p^c[S_p] \oplus_p H_p^r[S_p]$$

and using (1.4), we have

$$\begin{aligned} \|f\|_{S_p[F]}^p &= (1 - \theta) \left\| \sum_n \int_{\partial_0} f_n(z)^* f_n(z) d\mu_0(z) \right\|_{p/2}^{p/2} \\ &\quad + \theta \left\| \sum_n \int_{\partial_1} f_n(z) f_n(z)^* d\mu_1(z) \right\|_{p/2}^{p/2}. \end{aligned}$$

Combining the preceding inequalities with the classical Jensen inequality, we get (recalling that $p \geq 2$)

$$\begin{aligned} \left(\sum_n \|\alpha x_n \beta\|_2^2 \right)^{1/2} &= \left(\sum_n \|g_n(\theta)\|_2^2 \right)^{1/2} \\ &\leq \left[(1-\theta) \int_{\partial_0} \sum_n \|g_n(z)\|_2^2 d\mu_0(z) + \theta \int_{\partial_1} \sum_n \|g_n(z)\|_2^2 d\mu_1(z) \right]^{1/2} \\ &\leq \|f\|_{S_p[F]} \leq 1. \end{aligned}$$

Therefore, by Lemma 3.2, we deduce $\|x\|_{S_p[C_q]} \leq 1$, and so Q is completely contractive.

Now let $x \in S_p[C_q]$ with $\|x\|_{S_p[C_q]} < 1$. We have to find $f \in S_p[F]$ such that $f(\theta) = x$ and $\|f\|_{S_p[F]} < 1$. This is easy by interpolation. Indeed, by (3.1) and (1.1)

$$S_p[C_q] = (S_p[C_p], S_p[R_p])_\theta.$$

Thus there is $f \in \mathcal{F}(S_p[C_p], S_p[R_p])$ such that $f(\theta) = x$ and $\|f\|_{\mathcal{F}(S_p[C_p], S_p[R_p])} < 1$. Using the notation in the first part of the proof, we have

$$\left\| \sum_n \int_{\partial_0} f_n(z)^* f_n(z) d\mu_0(z) \right\|_{p/2} \leq \|f|_{\partial_0}\|_{L_2(\mu_0; S_p[C_p])}^2$$

and a similar inequality for the right endpoint. It then follows that $f \in S_p[F]$ and

$$\|f\|_{S_p[F]} \leq \|f\|_{\mathcal{F}(S_p[C_p], S_p[R_p])} < 1.$$

Therefore, Q is a complete quotient map, and so C_q is completely isometric to the quotient space $F/\ker(Q)$. $\square$

We turn to the second approach to Theorem 3.1 via the real interpolation. Below we will be rather brief and refer to [JX3] for more details and for more “concrete” representations of C_q as a quotient of subspace of $C_p \oplus_p R_p$.

We start by recalling the real interpolation method (see [BL] for more details). Let (X_0, X_1) be a compatible couple of Banach spaces. Let $X_0 +_p X_1$ denote the quotient of $X_0 \oplus_p X_1$ by the subspace $\{(x_0, x_1) : x_0 + x_1 = 0\}$. Similarly, we denote by $X_0 \cap_p X_1$ the diagonal subspace of $X_0 \oplus_p X_1$. Then $(X_0, X_1)_{\theta, p; K}$ is defined as the subspace of

$$L_p(X_0; t^{-\theta}) +_p L_p(X_1; t^{1-\theta})$$

consisting of all constant functions. Here for a given space X and $\alpha \in \mathbb{R}$ we have denoted by $L_p(X; t^\alpha)$ the X -valued L_p -space over $\mathbb{R}_+$ with respect to the weighted measure $t^{p\alpha} dt/t$. $(X_0, X_1)_{\theta, p; K}$ can be also described by the J-method. Let $(X_0, X_1)_{\theta, p; J}$ be the quotient space of

$$L_p(X_0; t^{-\theta}) \cap_p L_p(X_1; t^{1-\theta})$$

by the subspace of all f such that $\int_0^\infty f(t) dt/t = 0$. Then $(X_0, X_1)_{\theta, p; K} = (X_0, X_1)_{\theta, p; J}$ isomorphically with relevant constants depending only on θ and p .

Now let

$$K_{\theta, p} = (L_2(\ell_2; t^{-\theta}))_p^c +_p (L_2(\ell_2; t^{1-\theta}))_p^r$$

(recall that $(L_2(\ell_2; t^\alpha))_p^c$ means the Hilbert space $L_2(\ell_2; t^\alpha)$ equipped with the p -column space structure). We equip $K_{\theta, p}$ with the quotient operator space structure given by that of $(L_2(\ell_2; t^{-\theta}))_p^c \oplus_p (L_2(\ell_2; t^{1-\theta}))_p^r$. Let $C_{\theta, p; K}$ be the subspace of $K_{\theta, p}$ consisting of all constant functions. Similarly, set

$$J_{\theta, p} = (L_2(\ell_2; t^{-\theta}))_p^c \cap_p (L_2(\ell_2; t^{1-\theta}))_p^r,$$

and let $C_{\theta,p;J}$ be the quotient space of $J_{\theta,p}$ by the subspace of mean zero functions. We refer to [X3] for more discussions on such spaces.

Note that $K_{\theta,p}$ (resp. $J_{\theta,p}$) is a quotient (resp. a subspace) of $C_p \oplus_p R_p$. Thus both $C_{\theta,p;K}$ and $C_{\theta,p;J}$ are quotients of subspace of $C_p \oplus_p R_p$.

Theorem 3.3 *Let $1 \leq p \leq \infty$, $\min(p, p') < q < \max(p, p')$ and θ be determined by $\frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{p'}$. Then $C_q = C_{\theta,p;K}$ and $C_q = C_{\theta,p;J}$ completely isomorphically with relevant constants majorized by $2^{|1-\frac{2}{p}|}$.*

Proof. We first show the complete inclusion $C_{\theta,p;K} \subset C_q$ in the case of $p \geq 2$. Fix an $n \in \mathbb{N}$ and let $x = \sum_k x_k \otimes e_k \in S_p^n[C_{\theta,p;K}]$ of norm < 1 . (As before, we use (e_k) to denote the canonical basis of ℓ_2 .) Then there are $f \in S_p^n[(L_2(\ell_2; t^{-\theta}))_p^c]$ and $g \in S_p^n[(L_2(\ell_2; t^{1-\theta}))_p^r]$ such that $x = f(t) + g(t)$ for almost all $t \in \mathbb{R}$ and

$$\left(\|f\|_{S_p^n[(L_2(\ell_2; t^{-\theta}))_p^c]}^p + \|g\|_{S_p^n[(L_2(\ell_2; t^{1-\theta}))_p^r]}^p \right)^{\frac{1}{p}} < 1.$$

Using (1.4), we have

$$\begin{aligned} \|f\|_{S_p^n[(L_2(\ell_2; t^{-\theta}))_p^c]} &= \left\| \int_0^\infty \sum_k f_k(t)^* f_k(t) t^{-2\theta} \frac{dt}{t} \right\|_{S_{p/2}^n}^{\frac{1}{2}}, \\ \|g\|_{S_p^n[(L_2(\ell_2; t^{1-\theta}))_p^r]} &= \left\| \int_0^\infty \sum_k g_k(t) g_k(t)^* t^{2(1-\theta)} \frac{dt}{t} \right\|_{S_{p/2}^n}^{\frac{1}{2}}. \end{aligned}$$

To estimate the norm $\|x\|_{S_p^n[C_q]}$, we use again Lemma 3.2. Let α (resp. β) be a positive unit element in $S_{2r\theta-1}^n$ (resp. $S_{2r(1-\theta)-1}^n$). By perturbation, we can assume α and β invertible. Let A_0 be the right multiplication operator on $H = \ell_2(S_2^n)$ by $\beta^{\frac{1}{1-\theta}}$: $(x_k)_k \mapsto (x_k \beta^{\frac{1}{1-\theta}})_k$. Similarly, let A_1 be the left multiplication operator by $\alpha^{\frac{1}{\theta}}$. Then A_0 and A_1 are commuting invertible positive bounded operators on H . A_i induces an equivalent norm on H : $\|x\|_i = \|A_i(x)\|$ ($i = 0, 1$). Let H_i be H equipped with $\|\cdot\|_i$. Then (H_0, H_1) becomes a compatible couple of Hilbert spaces, which can be identified as a compatible couple of weighted L_2 -spaces. Using the real interpolation for weighted L_2 -spaces (see [BL]), we easily get the following:

$$\|A_0^{1-\theta} A_1^\theta(x)\| = \|x\|_{(H_0, H_1)_{\theta, 2; K}}, \quad x \in H.$$

Namely,

$$\left(\sum_k \|\alpha x_k \beta\|_2^2 \right)^{\frac{1}{2}} = \|x\|_{(H_0, H_1)_{\theta, 2; K}}.$$

The last interpolation norm is estimated as follows. First,

$$\begin{aligned} \|f\|_{L_2(H_0; t^{-\theta})}^2 &= \int_0^\infty \sum_k \|f_k(t)\|_{H_0}^2 t^{-2\theta} \frac{dt}{t} \\ &= \int_0^\infty \sum_k \text{Tr}(\beta^{\frac{1}{1-\theta}} f_k(t)^* f_k(t) \beta^{\frac{1}{1-\theta}}) t^{-2\theta} \frac{dt}{t} \\ &= \text{Tr}(\beta^{\frac{2}{1-\theta}} \int_0^\infty \sum_k f_k(t)^* f_k(t) t^{-2\theta} \frac{dt}{t}) \\ &\leq \left\| \int_0^\infty \sum_k f_k(t)^* f_k(t) t^{-2\theta} \frac{dt}{t} \right\|_{S_{p/2}^n}. \end{aligned}$$

Similarly,

$$\|g\|_{L_2(H_1; t^{1-\theta})}^2 \leq \left\| \int_0^\infty \sum_k g_k(t) g_k(t)^* t^{2(1-\theta)} \frac{dt}{t} \right\|_{S_{p/2}^n}.$$

Thus

$$\begin{aligned} (\|f\|_{L_2(H_0; t^{-\theta})}^2 + \|g\|_{L_2(H_1; t^{1-\theta})}^2)^{\frac{1}{2}} &\leq 2^{\frac{1}{2}-\frac{1}{p}} (\|f\|_{S_p^n[(L_2(\ell_2; t^{-\theta}))_p^c]}^p + \|g\|_{S_p^n[(L_2(\ell_2; t^{1-\theta}))_p^r]}^p)^{\frac{1}{p}} \\ &< 2^{\frac{1}{2}-\frac{1}{p}} \end{aligned}$$

Since $x = f(t) + g(t)$ for almost all $t \in \mathbb{R}_+$, we deduce that

$$\|x\|_{(H_0, H_1)_{\theta, 2, K}} < 2^{\frac{1}{2}-\frac{1}{p}};$$

whence

$$\left(\sum_k \|\alpha x_k \beta\|_2^2 \right)^{\frac{1}{2}} < 2^{\frac{1}{2}-\frac{1}{p}},$$

which, together with Lemma 3.2, implies that $x \in S_p^n[C_q]$ and $\|x\|_{S_p[C_q]} \leq 2^{\frac{1}{2}-\frac{1}{p}}$. Therefore, we deduce the complete inclusion $C_{\theta, p; K} \subset C_q$ and that its c.b. norm is majorized by $2^{\frac{1}{2}-\frac{1}{p}}$ in the case of $p \geq 2$.

To treat the case of $p \leq 2$ we recall the fact that for a Hilbert space H we have

$$(\overline{H_{p'}})^* \cong H_p^c \cong H_{p'}^r \quad \text{completely isometrically,}$$

where $\overline{H}$ denotes the conjugate of H . Then using the usual duality between sum and intersection spaces we obtain the following complete isometric identifications:

$$[(\overline{L_2(\ell_2; t^\theta)})_{p'}^c \cap_{p'} (\overline{L_2(\ell_2; t^{-(1-\theta)})})_{p'}^r]^* \cong K_{\theta, p}$$

and

$$\begin{aligned} [(\overline{L_2(\ell_2; t^\theta)})_{p'}^c \cap_{p'} (\overline{L_2(\ell_2; t^{-(1-\theta)})})_{p'}^r]^* &\cong (L_2(\ell_2; t^{-\theta}))_{p'}^r +_p (L_2(\ell_2; t^{(1-\theta)}))_{p'}^c \\ &\cong (L_2(\ell_2; t^{-(1-\theta)}))_{p'}^c +_p (L_2(\ell_2; t^\theta))_{p'}^r. \end{aligned}$$

However, since $p \leq 2$,

$$(L_2(\ell_2; t^{-(1-\theta)}))_{p'}^c +_p (L_2(\ell_2; t^\theta))_{p'}^r \subset K_{1-\theta, p'} \quad \text{completely contractively.}$$

Therefore, we deduce that

$$C_{\theta, p; K} \subset C_{1-\theta, p'; K} \quad \text{completely contractively.}$$

Applying the first part to $C_{1-\theta, p'; K}$ (with $p' \geq 2$), we get that $C_{1-\theta, p'; K} \subset C_q$ and the c.b. norm of this inclusion is $\leq 2^{\frac{1}{2}-\frac{1}{p'}}$, and so $C_{\theta, p; K} \subset C_q$ and is of c.b. norm $\leq 2^{\frac{1}{2}-\frac{1}{p}}$.

We have thus proved the complete inclusion $C_{\theta, p; K} \subset C_q$ for any $1 \leq p \leq \infty$ and its c.b. norm is majorized by $2^{|\frac{1}{2}-\frac{1}{p}|}$.

To show the inverse inclusion we use the J-method. In a similar way, we prove that $C_{\theta, p; J} \subset C_q$ and is of c.b. norm $\leq 2^{|\frac{1}{2}-\frac{1}{p}|}$. However, by the usual duality between the K- and J-methods, we easily check that

$$[C_{\theta, p; J}]^* = C_{1-\theta, p'; K} \quad \text{and} \quad [C_{\theta, p; K}]^* = C_{1-\theta, p'; J} \quad \text{completely isometrically.}$$

It then follows that $C_q = C_{\theta, p; K}$ and $C_q = C_{\theta, p; J}$ completely isomorphically with relevant constants bounded by $2^{|\frac{1}{2}-\frac{2}{p}|}$. $\square$

4 Noncommutative Khintchine inequalities

This section contains the main ingredient for the embedding of the quotients of subspace of $C_p \oplus_p R_p$ into noncommutative L_p -spaces. It is the noncommutative Khintchine type inequalities for Shlyakhtenko's generalized circular systems.

Given a complex Hilbert space H we denote as usual by $\mathcal{F}(H)$ the associated free Fock space:

$$\mathcal{F}(H) = \bigoplus_{n=0}^{\infty} H^{\otimes n},$$

where $H^{\otimes 0} = \mathbb{C}\Omega$ with Ω a unit vector, called the vacuum. Let $\ell(e)$ (resp. $\ell^*(e)$) denote the left creation (resp. annihilation) operator associated with a vector $e \in H$. Recall that $\ell^*(e) = (\ell(e))^*$.

Assume H is infinite dimensional and separable. Fix an orthonormal basis $\{e_{\pm k}\}_{k \geq 1}$ in H . We will also fix a sequence $\{\lambda_k\}_{k \geq 1}$ of positive numbers. Let

$$g_k = \ell(e_k) + \sqrt{\lambda_k} \ell^*(e_{-k}).$$

$(g_k)_{k \geq 1}$ is a generalized circular system in Shlyakhtenko's sense [S]. Let Γ be the von Neumann algebra on $\mathcal{F}(H)$ generated by the g_k . Let ρ be the vector state on Γ determined by Ω . By [S], ρ is faithful on Γ . By the identification of $L_1(\Gamma)$ with Γ_* , ρ is a positive unit element in $L_1(\Gamma)$, so for any $1 \leq p \leq \infty$, $\rho^{1/p}$ is a positive unit element in $L_p(\Gamma)$, and thus $g_k \rho^{1/p} \in L_p(\Gamma)$ for any k .

The following is the noncommutative Khintchine type inequalities for generalized circular systems from [X1]. See also [JX4] for this kind of inequalities in a more general setting. Note that the case $p = \infty$ is already contained in [PS].

Theorem 4.1 *Keep the notations above. Let M be a von Neumann algebra and (x_n) a finite sequence in $L_p(M)$.*

i) *If $p \geq 2$,*

$$\begin{aligned} & \max \left\{ \left\| \left(\sum_k x_k^* x_k \right)^{\frac{1}{2}} \right\|_{L_p(M)}, \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} x_k x_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\} \\ & \leq \left\| \sum_k x_k \otimes g_k \rho^{\frac{1}{p}} \right\|_{L_p(M \bar{\otimes} \Gamma)} \leq \\ & B_p \max \left\{ \left\| \left(\sum_k x_k^* x_k \right)^{\frac{1}{2}} \right\|_{L_p(M)}, \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} x_k x_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\}. \end{aligned}$$

ii) *If $p < 2$,*

$$\begin{aligned} & A_p^{-1} \inf \left\{ \left\| \left(\sum_k a_k^* a_k \right)^{\frac{1}{2}} \right\|_{L_p(M)} + \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} b_k b_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\} \\ & \leq \left\| \sum_k x_k \otimes g_k \rho^{\frac{1}{p}} \right\|_{L_p(M \bar{\otimes} \Gamma)} \leq \\ & \inf \left\{ \left\| \left(\sum_k a_k^* a_k \right)^{\frac{1}{2}} \right\|_{L_p(M)} + \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} b_k b_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\}, \end{aligned}$$

where the infimum runs over all decompositions $x_k = a_k + b_k$ in $L_p(M)$. The two positive constants A_p and B_p depend only on p and can be controlled by a universal constant.

iii) *Let G_p be the closed subspace of $L_p(\Gamma)$ generated by $\{g_k \rho^{1/p}\}_{k \geq 1}$. Then G_p is completely complemented in $L_p(\Gamma)$ with constant $\leq 2^{|1-\frac{2}{p}|}$.*

By [S] the algebra Γ generated by the g_k is a type III_λ factor ($0 < \lambda \leq 1$) (except the case where all λ_k are equal to 1, which is the usual case of Voiculescu's circular systems). Γ is never hyperfinite. Recall that Γ is the free analogue of the classical Araki-Woods quasi-free CAR factors. The latter factors are hyperfinite type III_λ . Thus it is natural to ask whether similar Khintchine inequalities also hold in the classical quasi-free case. This is indeed true for $1 < p < \infty$. To state this result, we need to introduce the Fermionic analogue of the system (g_k) .

Again consider a complex Hilbert space H as before. We denote by $\mathcal{F}_{-1}(H)$ the associated antisymmetric Fock space:

$$\mathcal{F}_{-1}(H) = \bigoplus_{n=0}^{\infty} \Lambda^n(H),$$

where $\Lambda^n(H)$ stands for the n -fold antisymmetric tensor product of H (with $\Lambda^0(H) = \mathbb{C}\Omega$). Let $c(e)$ (resp. $c^*(e)$) denote the creation (resp. annihilation) operator associated with a vector $e \in H$. Recall that $c^*(e) = (c(e))^*$.

Let $\{e_{\pm k}\}_{k \geq 1}$ and $\{\lambda_k\}_{k \geq 1}$ be as before. Put

$$f_k = c(e_k) + \sqrt{\lambda_k} c^*(e_{-k}).$$

Then $(f_k)_{k \geq 1}$ is a CAR sequence. Let Γ_{-1} be the von Neumann algebra on $\mathcal{F}_{-1}(H)$ generated by the f_k and ρ_{-1} the vector state on Γ_{-1} determined by Ω . ρ_{-1} is a quasi-free state, which is faithful on Γ_{-1} .

The following is the Fermionic analogue of Theorem 4.1.

Theorem 4.2 *Let M be a von Neumann algebra, $1 < p < \infty$ and $(x_k) \subset L_p(M)$ a finite sequence.*

i) *If $2 \leq p < \infty$,*

$$\begin{aligned} & \max \left\{ \left\| \left(\sum_k x_k^* x_k \right)^{\frac{1}{2}} \right\|_{L_p(M)}, \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} x_k x_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\} \\ & \leq \left\| \sum_k x_k \otimes f_k \rho_{-1}^{\frac{1}{p}} \right\|_{L_p(M \bar{\otimes} \Gamma_{-1})} \leq \\ & B'_p \max \left\{ \left\| \left(\sum_k x_k^* x_k \right)^{\frac{1}{2}} \right\|_{L_p(M)}, \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} x_k x_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\}. \end{aligned}$$

ii) *If $1 < p < 2$,*

$$\begin{aligned} & A'_p \inf \left\{ \left\| \left(\sum_k a_k^* a_k \right)^{\frac{1}{2}} \right\|_{L_p(M)} + \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} b_k b_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\} \\ & \leq \left\| \sum_k x_k \otimes f_k \rho_{-1}^{\frac{1}{p}} \right\|_{L_p(M \bar{\otimes} \Gamma_{-1})} \leq \\ & \inf \left\{ \left\| \left(\sum_k a_k^* a_k \right)^{\frac{1}{2}} \right\|_{L_p(M)} + \left\| \left(\sum_k \lambda_k^{1-\frac{2}{p}} b_k b_k^* \right)^{\frac{1}{2}} \right\|_{L_p(M)} \right\}, \end{aligned}$$

where the infimum runs over all decompositions $x_k = a_k + b_k$ with $a_k, b_k \in L_p(M)$. Here the positive constants A'_p and B'_p depend only on p .

iii) *Let F_p be the closed subspace of $L_p(\Gamma_{-1})$ generated by $\{f_k \rho_{-1}^{\frac{1}{p}}\}_k$. Then F_p is completely complemented in $L_p(\Gamma_{-1})$.*

Proof. Part i) follows from the noncommutative Burkholder inequality from [JX1] (see also [JX2]). Parts ii) and iii) are proved in a similar way as their free counterparts in Theorem 4.1 (see [X1]). We omit the details. $\square$

Remark. Very recently, Junge [J4] proved that Theorem 4.2, iii) holds for $p = 1$ too. However, it seems still unknown whether the constant A'_p there can be controlled by an absolute constant.

5 Embeddings

This section is devoted to the problem of embedding C_q (and more generally the quotients of subspace of $C_p \oplus_p R_p$) into noncommutative L_p -spaces for $1 \leq p < q \leq 2$.

Theorem 5.1 *Let H and K be two Hilbert spaces. Let $1 \leq p < 2$ and X be a quotient of subspace of $H_p^c \oplus_p K_p^r$. Then there is a QWEP von Neumann algebra M such that X embeds completely isomorphically into $L_p(M)$. Moreover, the relevant constant can be majorized by a universal constant.*

If $p > 1$, M can be chosen hyperfinite.

Proof. For simplicity, we assume $H = K = \ell_2$. Let X be a quotient of subspace of $C_p \oplus_p R_p$. Then $Y = X^*$ is a quotient of subspace of $C_{p'} \oplus_{p'} R_{p'}$, so there is a subspace $E \subset C_{p'} \oplus_{p'} R_{p'}$ such that Y is a quotient of E . By Corollary 2.3, E is decomposed into a direct sum of three spaces: a p' -column space, a p' -row space and the graph of an injective closed densely defined operator with dense range from a p' -column space into a p' -row space. Clearly, we need only to consider the third term. Thus without loss of generality, assume $E = G(\Delta) \subset C_{p'} \oplus_{p'} R_{p'}$, where $\Delta : \ell_2 \rightarrow \ell_2$ is a positive injective closed densely defined operator with dense range (see Remark 2.2). By discretization, we may as well assume Δ diagonal: $\Delta(e_k) = \delta_k e_k$ for all k , where (e_k) is the canonical basis of ℓ_2 and (δ_k) a sequence of positive numbers. Note that E is simply the diagonal subspace of $C_{p'} \oplus_{p'} R_{p'}$ but with the second space weighted by the δ_k . Passing to duals, we deduce that E^* is the quotient of $C_p \oplus_p R_p$ by the subspace $\{(x, -x) : x \in \ell_2\}$, i.e. $E^* = C_p \oplus_p R_p$.

Now we are in a position to apply Theorem 4.1. Let λ_k be such that $\delta_k = \lambda_k^{1/2-1/p}$. Then by the above discussion and Theorem 4.1, ii), we see that E^* is completely isomorphic to G_p in Theorem 4.1. Therefore, $X = X^{**} = Y^*$ is completely isomorphic to a subspace of $G_p \subset L_p(\Gamma)$. By [PS], Γ is QWEP.

If $p > 1$, instead of Theorem 4.1, we can use Theorem 4.2. Then E^* is completely isomorphic to F_p in Theorem 4.2. □

Remark 5.2 i) By Theorem 4.1, the space G_p is completely complemented in $L_p(\Gamma)$. Thus the proof above shows that the graph of an operator from C_p into R_p is completely isomorphic to a completely complemented subspace of $L_p(\Gamma)$. This fact is useful for applications.

ii) By [J4], the von Neumann algebra M in Theorem 5.1 can be chosen to be hyperfinite in the case of $p = 1$ too.

Let $QS(C_p, R_p)$ denote the family of all operator spaces which are completely isomorphic to quotients of subspace of $H_p^c \oplus_p K_p^r$ for some Hilbert spaces H and K . It is clear that if X belongs to $QS(C_p, R_p)$, then so does X^* . It was proved in [X1] that if both X and X^* embed completely isomorphically into some $L_p(M)$ ($1 < p < 2$), then $X \in QS(C_p, R_p)$. This together with the previous theorem implies the following

Corollary 5.3 *Let $1 < p < 2$ and X be an operator space. Then $X \in QS(C_p, R_p)$ iff both X and X^* embed completely isomorphically into a noncommutative L_p -space.*

Remark. It was shown in [PS] that if X is exact and both X and X^* completely embed into an L_1 , then $X \in QS(C_1, R_1)$.

Theorem 5.4 *Let $1 \leq p < q \leq 2$. Then there is a QWEP type III factor M such that C_q embeds completely isomorphically into $L_p(M)$.*

The same assertion holds for R_q too.

Proof. By Theorem 3.1, C_q is a quotient of subspace of $C_p \oplus_p R_p$. Thus Theorem 5.1 implies that C_q embeds into an $L_p(M)$. Let us show that M can be taken to be a type III factor. Let E be the space introduced during the proof of Theorem 3.1. Since an analytic function on the strip S is uniquely determined by its values on ∂_0 (or ∂_1), we see that E is a graph. Thus by Remark 5.2, E embeds into $L_p(\Gamma)$. By [S], Γ is a type III factor. □

Remark 5.5 The factor Γ above is of type III_λ for some $\lambda \in (0, 1]$. Using Theorem 3.3 and the precise form of the space $K_{\theta,p}$ introduced there, one easily shows that for any $\lambda \in (0, 1]$, Γ can be chosen to be of type III_λ . This follows from an appropriate discretization of $K_{\theta,p}$ and the classification theorem in [S].

Pisier [P5] proved that C_q ($1 < q \leq 2$) cannot embed completely isomorphically into the predual of a semifinite von Neumann algebra. We now extend this result to our general setting.

Theorem 5.6 *Let $1 \leq p < q \leq 2$. Then neither C_q nor R_q embeds completely isomorphically into a noncommutative $L_p(M)$ with a semifinite von Neumann algebra M .*

Junge [J3] proved that with the same p, q as above, the Schatten class S_q embeds isomorphically into $L_p(\mathcal{R})$ at the Banach space level, where $\mathcal{R}$ is the hyperfinite factor of type II_1 . It was left open there whether one can do this at the operator space level. Theorem 5.6 clearly implies that this is impossible. Thus we have the

Corollary 5.7 *Let $1 \leq p < q \leq 2$. Then S_q cannot embed completely isomorphically into a semifinite noncommutative L_p -space.*

Remark. It is still unknown whether S_q embeds completely isomorphically into a type III non-commutative L_p .

For the proof of Theorem 5.6 we will need the following factorization of c.b. maps from $L_p(M)$ to C_q .

Lemma 5.8 *Let M be a von Neumann algebra and H a Hilbert space. Let $2 \leq p \leq \infty$, $0 \leq \theta \leq 1$ and $\frac{1}{q} = \frac{1-\theta}{p} + \frac{\theta}{p'}$. Then for any map $u : L_p(M) \rightarrow H_q^c$ the following assertions are equivalent:*

- i) u is c.b..
- ii) *There are two positive unit functionals $f, g \in (L_{p/2}(M))^*$ such that*

$$(5.1) \quad \|u(x)\| \leq c (f(x^*x))^{(1-\theta)/2} (g(xx^*))^{\theta/2}, \quad x \in L_p(M).$$

Moreover, if c denotes the best constant in (5.1), then

$$c_{p,\theta}^{-1} c \leq \|u\|_{cb} \leq c,$$

where $c_{p,\theta}$ is a positive constant depending only on p and θ , which can be controlled by an absolute constant.

This is proved in [P4] for $p = \infty$ and [X1] for $p < \infty$. Moreover, the same result holds for any u defined on a subspace E of $L_p(M)$ (E must be then assumed exact in the case $p = \infty$).

We will also need the following simple description of c.b. maps between C_p and C_q . This is probably known to experts.

Lemma 5.9 *Let $1 \leq p, q \leq \infty$ with $p \neq q$. Then*

$$CB(C_p, C_q) = S_{\frac{2pq}{|p-q|}} \quad \text{isometrically.}$$

Proof. We first consider the case where $p = \infty$. We have

$$CB(C, C_q) = R \otimes_{\min} C_q = (R \otimes_{\min} C, R \otimes_{\min} R)_{1/q} = (S_\infty, S_2)_{1/q} = S_{2q}.$$

Now assume $p > q$. Let $u \in CB(C_p, C_q)$. For any $v \in CB(C, C_p)$, $uv \in CB(C, C_q)$ and

$$\|uv\|_{CB(C, C_q)} \leq \|u\|_{CB(C_p, C_q)} \|v\|_{CB(C, C_p)}.$$

Thus

$$\|uv\|_{S_{2q}} \leq \|u\|_{CB(C_p, C_q)} \|v\|_{S_{2p}}.$$

Taking the supremum over all v in the unit ball of S_{2p} , we get

$$\|u\|_{S_{\frac{2pq}{p-q}}} \leq \|u\|_{CB(C_p, C_q)}.$$

To prove the converse inequality we use interpolation. Let $\theta = q/p$. Then

$$C_p = (C, C_q)_\theta.$$

Thus by the case above and interpolation

$$\begin{aligned} S_{\frac{2pq}{p-q}} &= (S_{2q}, B(\ell_2))_\theta = (CB(C, C_q), CB(C_q, C_q))_\theta \\ &\subset CB((C, C_q)_\theta, C_q) = CB(C_p, C_q), \end{aligned}$$

where the inclusion is completely contractive. Therefore, we have obtained

$$CB(C_p, C_q) = S_{\frac{2pq}{p-q}} \quad \text{for } p > q.$$

Passing to adjoints, we deduce the identity in the case of $p < q$. □

Remark. The lemma above immediately yields the following isometric identities:

$$CB(R_p, R_q) = S_{\frac{2pq}{|p-q|}} \quad \text{and} \quad CB(R_p, C_q) = S_{\frac{2pq}{|p+q-pq|}} = CB(C_p, R_q).$$

Proof of Theorem 5.6. Assume C_q is completely isomorphic to a subspace of $L_p(M)$, where M is a semifinite von Neumann algebra equipped with a normal semifinite faithful trace τ . For simplicity, assume C_q itself is a subspace of $L_p(M)$. Let $J : C_q \rightarrow L_p(M)$ be the inclusion map. Then $u \stackrel{\text{def}}{=} J^* : L_{p'}(M) \rightarrow C_{q'}$ is completely contractive. Recall that $1 \leq p < q \leq 2$. Thus p' and q' satisfy the assumption of Lemma 5.8. Therefore, by that lemma, there are positive functionals f and g in the unit ball of $(L_{p'/2}(M))^*$ such that (5.1) holds. Replacing f and g by $f + g$, we can assume $f = g$. Thus we have

$$\|u(x)\| \leq c (f(x^*x))^{(1-\theta)/2} (f(xx^*))^{\theta/2}, \quad x \in L_{p'}(M).$$

If $p > 1$, f can be viewed as a positive unit operator in $L_r(M)$, where r is the conjugate index of $p'/2$. On the other hand, if $p = 1$, f can be taken as a normal state on M for u is normal (see [P5] for more details). Thus in this case, f is again a positive unit operator in $L_1(M)$. Therefore, the above inequality can be rewritten as

$$(5.2) \quad \|u(x)\| \leq c (\tau(fx^*x))^{(1-\theta)/2} (\tau(fxx^*))^{\theta/2}, \quad x \in L_{p'}(M).$$

As in [P5], we are going to reduce M to a finite von Neumann algebra. To this end, given $\lambda > 1$ let e_λ be the spectral projection of f corresponding to the interval (λ^{-1}, λ) . Since $f \in L_r(M)$ with $r < \infty$, $\tau(e_\lambda) < \infty$ and $e_\lambda \rightarrow 1$ strongly as $\lambda \rightarrow \infty$. Decompose J as follows:

$$J = R_{e_\lambda} L_{e_\lambda} J + R_{e_\lambda} L_{e_\lambda^\perp} J + R_{e_\lambda^\perp} J \stackrel{\text{def}}{=} J_1 + J_2 + J_3,$$

where L_a (resp. R_a) denotes the left (resp. right) multiplication by a . Consequently,

$$u = uR_{e_\lambda} L_{e_\lambda} + uL_{e_\lambda} R_{e_\lambda^\perp} + uL_{e_\lambda^\perp} \stackrel{\text{def}}{=} u_1 + u_2 + u_3.$$

We show that $\|u_2\|_{cb}$ and $\|u_3\|_{cb}$ become very small for λ sufficiently large. By (5.2), for any $x \in L_{p'}(M)$

$$\begin{aligned}\|u_2(x)\|^2 &= \|u(e_\lambda x e_\lambda^\perp)\|^2 \leq c^2 (\tau(fe_\lambda^\perp x^* e_\lambda x e_\lambda^\perp))^{1-\theta} (\tau(fe_\lambda x e_\lambda^\perp x^* e_\lambda))^\theta \\ &\leq c^2 (\tau(fe_\lambda^\perp x^* x))^{1-\theta} (\tau(fxx^*))^\theta \\ &= c^2 \|fe_\lambda^\perp\|_r^{1-\theta} (\tau(gx^*x))^{1-\theta} (\tau(fxx^*))^\theta,\end{aligned}$$

where $g = \|fe_\lambda^\perp\|_r^{-1} fe_\lambda^\perp$ is a positive unit operator in $L_r(M)$. Thus applying Lemma 5.8 to u_2 , we get

$$\|u_2\|_{cb} \leq c' \|fe_\lambda^\perp\|_r^{(1-\theta)/2}.$$

Since $e_\lambda^\perp \rightarrow 0$ strongly as $\lambda \rightarrow \infty$, $\|fe_\lambda^\perp\|_r \rightarrow 0$. Therefore, $\|u_2\|_{cb}$ is small for large λ . Similarly, we get the the same assertion for u_3 . Passing to adjoints, we deduce that $\|J_2\|_{cb}$ and $\|J_3\|_{cb}$ are small for large λ . Consequently, J_1 is a perturbation of J , and so J_1 is a complete isomorphism (if λ is large enough). However, J_1 takes values in $L_{p'}(e_\lambda M e_\lambda)$ and the restriction of τ to $e_\lambda M e_\lambda$ is finite. On the other hand, fe_λ is bounded by λ . Therefore, replacing J by J_1 if necessary, we are reduced to the case where τ is finite, say, normalized, and where f is a positive bounded operator in M , say, the identity of M . Then (5.2) becomes,

$$(5.3) \quad \|u(x)\| \leq c (\tau(x^*x))^{1/2}, \quad x \in L_{p'}(M).$$

The rest of the proof is to show that J factors through a p -column space. For this we consider the scalar product $\langle y, x \rangle = \tau(y^*x)$ on $L_{p'}(M)$. Let H be the completion of $L_{p'}(M)$ with respect to this scalar product. It is clear that the identity on $L_{p'}(M)$ induces a contractive inclusion of $L_{p'}(M)$ into H , denoted by ι . On the other hand, (5.3) implies that there is a bounded operator $T : H \rightarrow C_{q'}$ such that $\|T\| \leq c$ and $u = T\iota$. We now equip H with the operator space structure of $H_{p'}^c$. Then ι is completely contractive. This was already proved in [X1]. We include a proof for completeness. By [P2, Lemma 1.7], it suffices to prove that $I_{S_{p'}} \otimes \iota$ extends to a contraction from $S_{p'}[L_{p'}(M)]$ to $S_{p'}[H_{p'}^c]$. Let $x = (x_{ij}) \in S_{p'}[L_{p'}(M)]$ be a finite matrix. Then

$$\|I_{S_{p'}} \otimes \iota(x)\|_{S_{p'}[H_{p'}^c]}^2 = \left\| \left(\tau \left(\sum_k x_{ki}^* x_{kj} \right) \right)_{ij} \right\|_{S_{p'/2}}.$$

Let α be a positive finite matrix in the unit ball of S_t (t being the conjugate index of $p'/2$). Then (with Tr the usual trace on $B(\ell_2)$)

$$\begin{aligned}\sum_{i,j} \alpha_{ij} \tau \left(\sum_k x_{ki}^* x_{kj} \right) &= \text{Tr} \otimes \tau ((\alpha \otimes 1) x^* x) \\ &\leq \|\alpha \otimes 1\|_{S_t[L_t(M)]} \|x^* x\|_{S_{p'/2}[L_{p'/2}(M)]} \leq \|x\|_{S_{p'}[L_{p'}(M)]}^2.\end{aligned}$$

Then taking the supremum over α , we obtain

$$\|I_{S_{p'}} \otimes \iota(x)\|_{S_{p'}[H_{p'}^c]}^2 \leq \|x\|_{S_{p'}[L_{p'}(M)]}^2.$$

Therefore, ι is completely contractive.

Returning back to the factorization $u = T\iota$ and passing to adjoints, we deduce that $J = \iota^* T^* \stackrel{\text{def}}{=} vw$. Hence the identity of C_q factors through $\bar{H}_p^c$ with $\|v\|_{cb} \leq 1$. Restricted to a subspace of $\bar{H}_p^c$ if necessary, v can be assumed to have values in $J(C_q) = C_q$. Then H must be separable and infinite dimensional. By Lemma 5.9, $v \in S_{\frac{2pq}{q-p}}$. Consequently, $J \in S_{\frac{2pq}{q-p}}$, a contradiction, which achieves the proof of Theorem 5.6. $\square$

Acknowledgements. We are very grateful to Eric Ricard for fruitful discussions.

References

- [BL] Bergh, J. and Löfström, J. *Interpolation spaces*. Springer-Verlag, Berlin, 1976.
- [ER] Effros, Ed. and Ruan, Z.-J. *Operator spaces*, volume 23 of *London Mathematical Society Monographs. New Series*. The Clarendon Press Oxford University Press, New York, 2000.
- [H] Haagerup, U. L^p -spaces associated with an arbitrary von Neumann algebra. In *Algèbres d'opérateurs et leurs applications en physique mathématique (Proc. Colloq., Marseille, 1977)*, volume 274 of *Colloq. Internat. CNRS*, pages 175–184. CNRS, Paris, 1979.
- [J1] Junge, M. Embedding of the operator space OH and the logarithmic ‘little Grothendieck inequality’. To appear in *Invent. Math.*
- [J2] Junge, M. Embedding of OH in the predual of the hyperfinite III_1 von Neumann algebra. In preparation.
- [J3] Junge, M. Embeddings of non-commutative L_p -spaces into non-commutative L_1 -spaces, $1 < p < 2$. *Geom. Funct. Anal.*, 10:389–406, 2000.
- [J4] Junge, M. Personal communication.
- [JX1] Junge, M. and Xu, Q. Noncommutative Burkholder/Rosenthal inequalities. *Ann. Probab.*, 31(2):948–995, 2003.
- [JX2] Junge, M. and Xu, Q. Noncommutative Burkholder/Rosenthal inequalities: Applications. To appear.
- [JX3] Junge, M. and Xu, Q. Quantum probabilistic tools in operator spaces. In preparation.
- [JX4] Junge, M. and Xu, Q. Free random series. In preparation.
- [Ko] Kosaki, H. Applications of the complex interpolation method to a von Neumann algebra: noncommutative L^p -spaces. *J. Funct. Anal.*, 56(1):29–78, 1984.
- [P1] Pisier, G. *Introduction to operator space theory*, volume 294 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2003.
- [P2] Pisier, G. Non-commutative vector valued L_p -spaces and completely p -summing maps. *Astérisque*, (247): 1998.
- [P3] Pisier, G. The operator Hilbert space OH , complex interpolation and tensor norms. *Mem. Amer. Math. Soc.*, 122(585): 1996.
- [P4] Pisier, G. Completely bounded maps into certain Hilbertian operator spaces. *Int. Math. Res. Not.*, 74:3983–4018, 2004.
- [P5] Pisier, G. Operator Hilbert space OH and type III von Neumann algebras. *Bull. London Math. Soc.*, 36:455–459, 2004.
- [PS] Pisier, G. and Shlyakhtenko, D. Grothendieck’s theorem for operator spaces. *Invent. Math.*, 150(1):185–217, 2002.
- [PW] Pusz, W. and Woronowicz, S. L. Functional calculus for sesquilinear forms and the purification map. *Rep. Mathematical Phys.*, 8: 159–170, 1975.
- [S] Shlyakhtenko, D. Free quasi-free states. *Pacific J. Math.*, 177(2):329–368, 1997.
- [Sp] Speicher, R. A new example of “independence” and “white noise”. *Probab. Theory Related Fields*, 84: 141–159, 1990.

- [Te1] Terp, M. L^p -spaces associated with von Neumann algebras. Notes, Math. Institute, Copenhagen Univ. 1981.
- [Te2] Terp, M. Interpolation spaces between a von Neumann algebra and its predual. *J. Operator Theory*, 8:327–360, 19827.
- [X1] Xu, Q. Operator space Grothendieck inequalities for noncommutative L_p -spaces. To appear.
- [X2] Xu, Q. A description of $(C_p[L_p(M)], R_p[L_p(M)])_\theta$. *Proc. Edinburgh Math. Soc.*, to appear.
- [X3] Xu, Q. Interpolation of operator spaces. *J. Funct. Anal.*, 139(2):500–539, 1996.